

## 6.2.1 Amines


### Naming Amines

These end in **-amine**.

There is, however, rather confusingly two ways of using this suffix.


The exam board tend to use the common version where the name stem ends in **-yl** **propylamine**.

The IUPAC version of the same chemical is **propan-1-amine**. (This is used in the same way as naming alcohols)



propylamine  
Or propan-1-amine


If there is another priority functional group as well as the amine group then the prefix **amino** is used.



2-aminopropanoic acid.

If the amine is secondary and has two alkyl groups attached to the nitrogen, then each chain is named and the smaller alkyl group is preceded by an **-N** which plays the same role as a number in positioning a side alkyl chain

$\text{CH}_3\text{CH}_2\text{CH}_2\text{NHCH}_3$   
N-methylpropylamine (common name)  
N-methylpropan-1-amine (IUPAC name)




In the common naming version if the chain lengths are the same an **-N** is not used

Diethylamine (common name- does not use N if chains are same length)

N-ethylethanamine (IUPAC name does still use N)

If a tertiary amine similar rules apply, and each alkyl side group is given an N


$\text{CH}_3\text{CH}_2\text{CH}_2-\text{N}(\text{CH}_3)_2$   
N,N-dimethylpropylamine (common name)  
N,N-dimethylpropan-1-amine (IUPAC name)




It could also be named  
1,6-diaminohexane

### Base Properties

Primary aliphatic amines act as Bronsted-Lowry Bases because the lone pair of electrons on the nitrogen is readily available for forming a dative covalent bond with a  $\text{H}^+$  and so accepting a proton.




Primary aliphatic amines are stronger bases than ammonia as the alkyl groups are electron releasing and push electrons towards the nitrogen atom and so make it a stronger base.




### Base strength of aromatic amines

Primary aromatic amines such as Phenylamine do not form basic solutions because the lone pair of electrons on the nitrogen delocalise with the ring of electrons in the benzene ring. This means the N is less able to accept protons.




### Reactions with acids

Amines as bases react with acids to form ammonium salts.



Addition of NaOH to an ammonium salt will convert it back to the amine




These ionic salts will be solid crystals, if the water is evaporated, because of the strong ionic interactions.

The ionic salts formed in this reaction means that the compounds are soluble in the acid.  
e.g. Phenylamine is not very soluble in water but phenylammonium chloride is soluble


## Nucleophilic properties

Primary amines can be formed by the **nucleophilic substitution** reaction between haloalkanes and ammonia.

### Reaction with ammonia forming primary amine




**Excess Ammonia dissolved in ethanol** is the reagent



Using an **excess of Ammonia** can limit further substitution reactions and will **maximise the amount of primary amine** formed

### Reacting primary amines with haloalkanes forming secondary amine


Amines will react with haloalkanes in the same nucleophilic substitution reactions that ammonia does above



A primary amine will react with a haloalkane to form a secondary amine

### Reacting secondary amines with haloalkanes to form a tertiary amine

The same reaction mechanism occurs with the secondary amine reacting to form a tertiary amine




### Reducing nitroarenes to aromatic amines

The nitro group on an arene can be reduced an amine group as follows

**Reagent:** Sn and concentrated HCl

**Conditions:** Heating

**Mechanism:** reduction



As the reaction is carried out in HCl the salt  $\text{C}_6\text{H}_5\text{NH}_3^+\text{Cl}^-$  will be formed. Reacting this salt with NaOH will give phenylamine.